



## Cold-water corals and large hydrozoans provide essential fish habitat for *Lappanella fasciata* and *Benthocometes robustus*



José Nuno Gomes-Pereira<sup>a,b,c,\*</sup>, Vanda Carmo<sup>a,h</sup>, Diana Catarino<sup>a</sup>, Joachim Jakobsen<sup>d</sup>, Helena Alvarez<sup>e</sup>, Ricardo Aguilar<sup>e</sup>, Justin Hart<sup>f</sup>, Eva Giacomello<sup>a</sup>, Gui Menezes<sup>a</sup>, Sergio Stefanni<sup>a,g</sup>, Ana Colaço<sup>a</sup>, Telmo Morato<sup>a</sup>, Ricardo S. Santos<sup>a</sup>, Fernando Tempera<sup>a</sup>, Filipe Porteiro<sup>a,h</sup>

<sup>a</sup> MARE – Marine and Environmental Sciences Centre, IMAR, Department of Oceanography and Fisheries, University of the Azores, 9901-862 Horta, Azores, Portugal

<sup>b</sup> Portuguese Task Group for the Extension of the Continental Shelf (EMEPC), Rua Costa Pinto 165, Paço de Arcos 2770-047, Portugal

<sup>c</sup> Naturalist, Science & Tourism, Rua da Ladeira n2, 9900-029 Horta, Faial Island, Portugal

<sup>d</sup> Rebikoff-Niggeler Foundation, Rocha Vermelha, Apt. 249, Praia do Almoxarife, PT-9900-909 Horta, Portugal

<sup>e</sup> Oceana, Gran Vía 59, 9º, 28013 Madrid, Spain

<sup>f</sup> Cetacean Watching Lda, Cais da Madalena, 9950-305 Madalena do Pico, Pico, Portugal

<sup>g</sup> Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Naples, Italy

<sup>h</sup> Direção Regional dos Assuntos do Mar, Secretaria Regional do Mar, Ciência e Tecnologia, Governo Regional dos Açores. 9901-014, Horta, Faial, Azores, Portugal

### ARTICLE INFO

#### Keywords:

Fish habitat  
Cold-water corals  
Hydrozoa  
*Benthocometes robustus*  
*Lappanella fasciata*  
Behavior  
Diet  
mtDNA COI  
Macaronesia  
Northeast Atlantic Ocean

### ABSTRACT

Many fish species are well-known obligatory inhabitants of shallow-water tropical coral reefs but such associations are difficult to study in deep-water environments. We address the association between two deep-sea fish with low mobility and large sessile invertebrates using a compilation of 20 years of unpublished *in situ* observations. Data were collected on Northeast Atlantic (NEA) island slopes and seamounts, from the Azores to the Canary Islands, comprising 127 new records of the circalittoral Labridae *Lappanella fasciata* and 15 of the upper bathyal Ophiididae *Benthocometes robustus*. Observations by divers, remote operated vehicles (ROV *SP*, *Luso*, *Victor*, *Falcon Seaeye*), towed vehicles (Greenpeace) and manned submersibles (*LULA*, *Nautile*) validated the species association to cold water corals (CWC) and large hydrozoans. *L. fasciata* occurred from lower infralittoral (41 m) throughout the circalittoral, down to the upper bathyal at 398 m depth. Smaller fishes (< 10 cm) tend to form larger schools up to five individuals, with larger fishes (10–15 cm) occurring alone or in smaller groups at greater depths. The labrids favoured areas with large sessile invertebrates (> 10 cm) occurring at < 1 body-length, swimming inside or in close vicinity to the tallest and most complex three-dimensional structure in the field of observation. These included hydrozoans (*Polyplumaria flabellata*, *Nemertesia antennina*), CWC (e.g. *Antipathella wollastoni*, *Acanthogorgia armata*, *Stichopathes* sp.), and less frequently sponges (e.g. *Pseudotrachya hystrix*). *B. robustus* presented a coral-cryptic behavior, being recorded in the bathyal zone between 350 and 734 m depth, always inside CWC (e.g. *Acanthogorgia* spp., *Antipathella* spp., *Callogorgia verticillata*, *Dendrophyllia alternata*, *Leiopathes* spp.), and remaining within the coral branching. *B. robustus* were collected with baited traps providing biological information and dietary information reinforcing the trophic linkage between the CWC habitat and this predator. Gathered evidence renders CWC and hydroid gardens as Essential Fish Habitats for both species, being therefore sensitive to environmental and anthropogenic impacts on these Vulnerable Marine Ecosystems. The Mediterranean distribution of *L. fasciata* is extended to NEA seamounts and island slopes and the ampho-Atlantic distribution of *B. robustus* is bridged with molecular data support. Both species are expected to occur throughout the Macaronesia and Mediterranean island slopes and shallow seamounts on habitats with large sessile invertebrates.

### 1. Introduction

Understanding fish habitats in the deep ocean has long been

recognized as a complex task (Benaka, 1999; Auster, 2005). Efforts in this direction have included both observations from traditional fishing and acoustic surveys (Collie et al., 1997; Kutti et al., 2014) as well as

\* Corresponding author at: MARE – Marine and Environmental Sciences Centre, IMAR, Department of Oceanography and Fisheries, University of the Azores, 9901-862 Horta, Azores, Portugal.

E-mail address: [contact@naturalist.pt](mailto:contact@naturalist.pt) (J.N. Gomes-Pereira).

<http://dx.doi.org/10.1016/j.dsr2.2017.09.015>